

Domination of black hole accretion in brane cosmology

A. S. Majumdar*

S. N. Bose National Centre for Basic Sciences, Block JD, Sector III, Salt Lake, Kolkata 700098, India

We consider the evolution of primordial black holes formed during the high energy phase of the braneworld scenario. We show that the effect of accretion from the surrounding radiation bath is dominant compared to evaporation for such black holes. This feature lasts till the onset of matter (or black hole) domination of the total energy density which could occur either in the high energy phase or later. We find that the black hole evaporation times could be significantly large even for black holes with small initial mass to survive till several cosmologically interesting eras.

PACS number(s): 98.80.Cq

The braneworld scenario in which our observable universe is a brane embedded in a higher dimensional bulk has gained much popularity in recent times. The motivation for such a consideration originates from solutions of string theory where matter and radiation are confined to the 3-brane, whereas gravity propagates also in the bulk. The significant consequences of this idea on the physics of the early universe is the focus of much current attention [1]. The foremost departure from standard cosmology is that there exists a regime during the early stages when the expansion rate of the universe is proportional to its energy density. The aim is to look for observable implications of this behaviour on cosmology.

Primordial black holes through their evolution and evaporation are potentially interesting candidates towards probing the high energy early stages of brane cosmology. The formation of primordial black holes through various mechanisms has been studied in much detail [2]. More recently, the diverse cosmological ramifications of primordial black holes are being vigorously pursued. Several papers have analysed the implications of a population of primordial black holes in the early universe on density perturbations, baryogenesis, reionization and nucleosynthesis [3,4]. Other investigations have focussed on methods of constraining the spectra of primordial black holes using observational results on gamma rays, background neutrinos, dark matter and quintessence [5]. Recently, several attempts of describing the formation, evaporation and cosmological consequences of primordial black holes in the braneworld scenario have also been undertaken [6,7].

In the present Letter, we consider the evolution of primordial black holes formed during the high energy brane phase when the expansion rate of the universe is linearly proportional to its energy density [1]. We restrict ourselves to the Randall-Sundrum Type II model [8]. Black holes formed during this regime evaporate at a rate proportional to their effective area times the fourth power of their temperature. Using the induced metric on the

brane, and expressions for the black hole radius and temperature in terms of the AdS radius l , Guedens et al [7] derived an evaporation law which is different from that of usual 4D black holes. However, for a more complete analysis of the evolution of black holes, one must take into account the effect of accretion from the surrounding radiation bath as well. This point has been emphasized by Zeldovich and Novikov [2] and the ramifications of considering accreting black holes have been found to be significant in the context of standard cosmology [4]. In our analysis, we consider the effect of both accretion and evaporation on primordial black holes formed on the brane.

Let us consider the rate of change of mass $\dot{M}$ of a black hole immersed in a radiation bath. The accretion rate is proportional to the surface area of the black hole times the energy density of radiation. The evaporation rate is proportional to the surface area times the fourth power of temperature. Thus, $\dot{M}$ is given by

$$\dot{M} = 4\pi r_{BH}^2 \left(-g_{brane} \sigma T_{BH}^4 + \rho_R \right) \quad (1)$$

where g_{brane} represents the effective number of degrees of freedom on the brane, r_{BH} and T_{BH} are the radius and temperature of the black hole, σ is the Stefan-Boltzmann constant, and ρ_R is the energy density of radiation given by

$$\rho_R = \frac{3M_4^2}{32\pi t_c t} \quad (2)$$

$$r_{BH} = \left(\frac{8}{3\pi} \right)^{1/2} \left(\frac{l}{l_4} \right)^{1/2} \left(\frac{M}{M_4} \right)^{1/2} l_4 \quad (3)$$

$$T_{BH} = \frac{1}{2\pi r_{BH}} \quad (4)$$

$$\sigma = \frac{\Gamma(4)\zeta(4)}{8(\pi)^2} \quad (5)$$

*Electronic address: archan@bose.res.in

where M_4 and l_4 are the four dimensional Planck mass and length, respectively, ($t_c \equiv l/2$) is the transition time to standard cosmology, and the above relations hold for $t \ll t_c$. (Note here that we have neglected the contribution of evaporation into the bulk given by a term proportional to $4\pi r_{BH}^2 g_{bulk} T_{BH}^5$ since it is subdominant even for very small black holes [7]). With the above substitutions the black hole equation can be written as

$$\dot{M} = -\frac{AM_4^2}{Mt_c} + \frac{BM}{t} \quad (6)$$

where ($A \simeq 3/(16)^3\pi$) and ($B \simeq 2/\pi$) are dimensionless numbers¹. The exact solution for Eq.(6) is

$$M(t) = \left[\left(M_0^2 - \frac{2AM_4^2}{2B-1} \frac{t_0}{t_c} \right) \left(\frac{t}{t_0} \right)^{2B} + \frac{2AM_4^2}{2B-1} \frac{t}{t_c} \right]^{1/2} \quad (7)$$

with $M = M_0$ at the time of BH formation $t = t_0$. It can be seen that an extremum for the function $M(t)$ exists only if $(M_0/M_4)^2 < (2A/2B-1)(t_0/t_c)$. The relation between formation time and mass [7]

$$\frac{t_0}{t_4} \simeq \frac{1}{4} \left(\frac{M_0}{M_4} \right)^{1/2} \left(\frac{l}{l_4} \right)^{1/2} \quad (8)$$

shows that such a small initial mass will violate the lower bound on horizon mass ($M_H/M_4 > 2 \times 10^6 (l_4/l)^{1/3}$) obtained from the contribution of gravitational waves towards CMBR anisotropy [9]. This result shows that for a radiation dominated high energy phase ($t < t_c$), accretion dominates over evaporation for all times and the mass of a primordial black hole continues to grow monotonically with

$$\frac{M(t)}{M_0} \simeq \left(\frac{t}{t_0} \right)^B \quad (9)$$

Let us now consider the scenario in which a certain number density of primordial black holes exchange energy with the surrounding radiation by accretion and evaporation. Black holes are formed with an initial mass spectrum whose range is much debated [10]. In order to keep our analysis uncomplicated, the subsequent calculations are performed by considering an average initial mass M_0 . We assume that at a time t_0 the fraction of the total energy in black holes is β , and the number density of black holes is $n_{BH}(t_0)$. Hence, we have

$$\rho_T(t_0) = \rho_R(t_0) + \rho_{BH}(t_0) \quad (10)$$

$$\rho_{BH}(t_0) = \beta \rho_T(t_0) = M_0 n_{BH}(t_0) \quad (11)$$

$$\rho_R(t_0) = (1 - \beta) \rho_T(t_0) \quad (12)$$

The number density of black holes $n_{BH}(t)$ scales as $a(t)^{-3}$, and thus for a radiation dominated evolution on the brane, one has $(n_{BH}(t)/n_{BH}(t_0)) = (t_0/t)^{3/4}$ since $a(t) \propto t^{1/4}$. The net energy in black holes grows since accretion dominates over evaporation. The condition for the universe to remain radiation dominated (i.e., $\rho_{BH}(t) < \rho_R(t)$) at any instant t can be derived from Eqs.(7,10,11,12) to be

$$\beta < \frac{(t_0/t)^{B+1/4}}{1 + (t_0/t)^{B+1/4}} \quad (13)$$

Thus depending on the value of the parameter β there may or may not be an era of black hole domination in the early brane dominated case. We will analyse these two cases separately.

We first consider the situation when the cosmology stays radiation dominated up to the time when brane effects are important, i.e., $t \leq t_c$. For this to be the case, the initial fraction β should be such that

$$\frac{\beta}{1 - \beta} < \left(\frac{t_0}{t_c} \right)^{B+1/4} \quad (14)$$

If the universe remains radiation dominated up to time t_c , then (for $M_0 \geq M_4$) the mass of a black hole is given by Eq.(9). However, the black hole must remain small enough, i.e., $(M/M_4) < (3\pi/4)(t/t_4)$ for it to obey the 5D evaporation law that we have used [7]. These criteria can be used to put an upper bound on the initial mass:

$$\frac{M_0}{M_4} < \left(\frac{3\pi}{4(2\sqrt{2})^B} \right)^{\frac{2}{2-B}} \frac{t_c}{t_4} \quad (15)$$

In the low energy regime $\rho_R \propto t^{-2}$. The black hole mass will continue to grow due to accretion up to a certain time (say t_t) after which the radiation density becomes too dilute for accretion to be significant. At this stage the rate of evaporation is also insignificant since the black hole masses have grown by several orders of magnitude from their initial values. So there ensues an era during which the black hole mass stays nearly constant over a period of time until evaporation takes over finally. This feature is well supported by numerical simulations of coupled black hole and Einstein equations for 4D black

¹We assume that the black holes can emit massless particles only and take $g_{brane} = 7.25$ as in [7]. It is known that the cross section for particle absorption is increased for relativistic particles [2], and also for relativistic motion of black holes [4]. In the present analysis we neglect these effects which tend to increase the value of B by a small amount.

holes [4]. For the present case the accretion rate is smaller since the surface area is $\propto M$ instead of M^2 for $4D$ black holes. Furthermore, the evaporation rate is $(\propto M^{-1})$ instead of M^{-2} . Hence, a critical mass M_{max} is reached before evaporation starts dominating from the transition time t_t onwards². Assuming the small mass condition $((M/M_4) < (3\pi/4)(t/t_4))$ to be still valid together with radiation domination at the time t_t , the lifetime for a typical black hole can be computed from Eq.(6) (using $M = M_{max}$ at $t = t_t$) and is given by

$$\frac{t_{end}}{t_4} \simeq \frac{1}{2A} \left(\frac{M_{max}}{M_4} \right)^2 \frac{t_c}{t_4} \quad (16)$$

Using Eqs.(8) and (9), the lifetime is given by

$$\frac{t_{end}}{t_4} \simeq \frac{4}{A} (2\sqrt{2})^B \left(\frac{M_0}{M_4} \right)^{2-B} \frac{t_c}{t_4} \left(\frac{t_t^2}{t_c t_4} \right)^B \quad (17)$$

Guedens et al [7] derived the increase of black hole evaporation time compared to $4D$ black holes as $\propto (l/r_0)^2$, i.e., $\{[t_{end}(M_0, 5D)]/[t_{end}(M_0, 4D)] \simeq [(t_c/t_4)(M_4/M_0)]\}$. Our results show that over and above the enhancement of black hole lifetime originating from a modified evaporation law, the black hole lifetime is further increased by a factor $\propto (M_{max}/M_0)^2$ purely due to accretion given by

$$\frac{t_{end}(M, 5D)}{t_{end}(M_0, 5D)} \simeq \left(\frac{M_4}{M_0} \right)^B \left(\frac{t_t^2}{t_c t_4} \right)^B \quad (18)$$

for $t_c \leq t_t \leq t_{eq}$. We thus find that the effect of accretion in prolonging the life time of black holes is more pronounced compared to the $5D$ evaporation effect for small black holes. From Eq.(17) it can be seen that for $(l/l_4) \sim 10^{20}$, a small number density of primordial $5D$ black holes formed with very low or even subplanckian masses ($M_0 \leq M_4$) could survive up to the era of nucleosynthesis and beyond if $t_t \geq 10^5 t_c$. This result is likely to have multifarious cosmological consequences [3,5] which need to be investigated in detail. But for black holes with large initial masses ($M_0 \gg M_4$) which form later, there is a lower net mass gain. Hence, for example, if we take $t_t \sim 10^5 t_c$, and $(l/l_4) \sim 10^{20}$, black holes formed with $M_0 = 10^{15} M_4 \simeq 10^{10} \text{g}$, will have $M_{max} \simeq 10^{15} \text{g}$, and will be evaporating now. However, if the AdS radius is the maximum allowed by the current experimental bounds [11], i.e., $(l/l_4) \simeq 10^{30}$ black holes with initial masses as low as $M_0 = 10^8 M_4 \simeq 10^3 \text{g}$ survive up to the present era even for $t_t \sim t_c$.

We next consider the case when the initial energy fraction β is such that the accretion process causes $\rho_{BH}(t)$ to exceed $\rho_R(t)$ at a time $t < t_c$. If β exceeds the bound (we still assume $\beta \ll O(1)$) given in Eq.(13), then the cosmology enters a matter (black hole) dominated regime in the high energy phase when $H^2 \propto \rho^2$. The onset of such a matter dominated era is labelled by t_{heq} which can be written from Eqs.(9-12) as³

$$\frac{t_{heq}}{t_0} = \left(\frac{1-\beta}{\beta} \right)^{\frac{4}{4B+1}} \equiv \gamma \quad (19)$$

The mass of a black hole at t_{heq} is given by $M(t_{heq}/M_0) = \gamma^B$.

For $t > t_{heq}$ the Hubble expansion is essentially driven by the black holes ($p = 0$ and $H \propto \rho_{BH}$). The number density of black holes scales as matter ($n_{BH}(t) \propto a^{-3}$), and thus the scale factor grows as $a \sim t^{1/3}$. During this era, the radiation density ρ_R is governed by the equation

$$\frac{d}{dt} \left(\rho_R(t) a^4(t) \right) = -\dot{M}(t) n_{BH}(t) a(t) \quad (20)$$

where the r.h.s (contribution from accreting black holes) cannot be neglected in comparison with the redshifting term ($\rho_R \sim a^{-4}$). Taking into account the radiation dominated expansion for $t < t_{heq}$, and black hole dominated expansion for $t > t_{heq}$, one has

$$\frac{n_{BH}(t)}{n_{BH}(t_0)} = \left(\frac{t_0}{t_{heq}} \right)^{3/4} \frac{t_{heq}}{t} \quad (21)$$

Using Eqs.(9-12) and (21) in (20) gives

$$\dot{\rho}_R + \frac{4\rho_R}{3t} = -\beta B \gamma^{1/4} \frac{\rho(t_0)}{t_0} \left(\frac{t}{t_0} \right)^{B-2} \quad (22)$$

Using the condition of matter-radiation equality at t_{heq} , i.e., $[\rho_R(t_{heq}) = \rho_{BH}(t_{heq}) = M(t_{heq}) n_{BH}(t_{heq})]$, one can solve Eq.(22) to obtain

$$\rho_R(t) \approx \gamma^{-1} \rho(t_0) \left(\frac{t_{heq}}{t} \right) - \frac{\beta B}{B+1/3} \gamma^{1/4} \rho(t_0) \left(\frac{t_0}{t} \right)^{1-B} \quad (23)$$

where we have neglected a term of higher order in t_{heq}/t .

The black hole equation for $t > t_{heq}$ is given by

²The exact determination of t_t requires the numerical integration of the coupled Einstein and black hole equations (see, for example [4]) and is beyond the scope of the present paper.

³Although the formation time t_0 of individual black holes is spread out in accordance with the initial mass spectra, we have used t_0 to signify the mean formation time in the similar sense of using M_0 as the mean initial mass.

$$\dot{M} = B\gamma^{-1} \frac{M}{t_0} \left(\frac{t_{heq}}{t} \right) - C\gamma^{1/4} \frac{M}{t_0} \left(\frac{t_0}{t} \right)^{1-B} \quad (24)$$

where $C = \beta B / (B + 1/3)$. Note that we have ignored the evaporation term at this stage since it is even more negligible compared to accretion because of the resultant mass gain up to t_{heq} . Using $M(t_{heq}/M_0) = \gamma^B$, the solution of Eq.(24) can be obtained. Beyond t_{heq} the black holes continue to gain mass until the two terms in Eq.(24) become comparable. At this stage one enters the transition regime (t_t) after which evaporation takes over. Setting the r.h.s of Eq.(24) to 0 at $t = t_t$, one obtains

$$\frac{t_t}{t_{heq}} = \left(\frac{B + 1/3}{1 - \beta} \right)^{\frac{3}{3B+1}} \equiv \delta \quad (25)$$

Thus the accretion regime lasts for a brief duration in the matter dominated phase, and the maximum black hole mass $M(t_t)$ is given from the solution of Eq.(24) by

$$\begin{aligned} \frac{M(t_t)}{M(t_{heq})} = \exp & \left[3B \left(1 - \left(\frac{\beta}{B + 1/3} \right)^{\frac{1}{3B+1}} \right) \right. \\ & \left. + \frac{\beta}{B + 1/3} \left(1 - \left(\frac{B + 1/3}{\beta} \right)^{\frac{3B}{3B+1}} \right) \right] \end{aligned} \quad (26)$$

In the evaporating regime ($t > t_t$), the rate of change of black hole mass is given by the first term in Eq.(6). This can be integrated to give

$$M(t) = \left[M^2(t_t) - \frac{2AM_4^2}{t_c}(t - t_t) \right]^{1/2} \quad (27)$$

The energy density in radiation $\rho_R(t)$ whose time evolution is governed by Eq.(20) now has a positive contribution from black hole evaporation. Making the appropriate substitutions, one can again integrate $\dot{\rho}_R(t)$ to obtain

$$\rho_R(t) \approx 3\beta\rho(t_0)\gamma^{-3/4} \frac{M(t_t)}{M_0} \left[\delta^{-1} - \frac{t_{heq}}{t} \right] \quad (28)$$

The universe gets reheated as $\rho_R(t)$ increases with time. The stage of black hole domination lasts up to a time t_r when $\rho_R(t_r) = n_{BH}(t_r)M(t_r)$. Subsequently, radiation domination takes over again. From Eqs.(27) and (28),

$$\frac{t_c}{t_r} \approx \frac{3}{2\delta\gamma} + \frac{A}{4\gamma^{2B}} \left(\frac{M_4}{M_0} \right)^2 \quad (29)$$

The standard low energy cosmology should be radiation dominated with sufficient radiation temperature for nucleosynthesis to occur. Thus the era of black hole domination should be over clearly before nucleosynthesis. Since it is possible for t_c/t_4 to be as large as 10^{30} [11], let us simply demand that the onset of radiation

domination take place before t_c . Requiring $t_r < t_c$, one gets a lower bound on β from Eq.(29), i.e.,

$$\beta \geq \left[\frac{4t_0}{3t_c} (B + 1/3)^{\frac{3}{3B+1}} \right]^{B+1/4} \quad (30)$$

The evaporation time of black holes in this scenario can be calculated from Eq.(27). The mass gain between t_{heq} and t_t given by Eq.(26) is negligible if Eq.(30) is satisfied and also $\beta \ll O(1)$. The black hole lifetime is

$$\frac{t_{end}}{t_4} \approx \left(\frac{M_0}{M_4} \right)^2 \frac{t_c}{t_4} \gamma^{2B} \quad (31)$$

The contribution of accretion in prolonging the lifetime of black holes compared to the modified evaporation effect is given in this scenario by

$$\frac{t_{end}(M, 5D)}{t_{end}(M_0, 5D)} = \left(\frac{M_{max}}{M_0} \right)^2 \approx \gamma^{2B} \quad (32)$$

This enhancement is independent of the value of the initial mass. Thus in this case the lifetime of the whole mass spectrum gets prolonged due to accretion by the same scaling factor γ^{2B} , as distinct from the scenario of radiation domination throughout where the low mass black holes are affected most. Thus, for instance, if $l/l_4 \simeq 10^{20}$ and $\beta = 10^{-3}$, black holes with $M_0 \simeq 10^{12}g$ evaporate during the present era.

To summarize, we have studied the evolution of primordial black holes formed during the radiation dominated high energy phase on the brane. These black holes obey a modified evaporation law [7]. We have shown that the accretion of surrounding radiation completely dominates evaporation as long as radiation domination persists. This results in the net growth of mass of the black holes. Compared to the case of black holes formed with the same initial mass in standard cosmology, the black holes in the braneworld scenario evaporate much later. We have found that the effect of accretion in enhancement of black hole lifetime could be very significant for even low mass black holes for a wide range of parameters. It needs to be emphasized that the effect of accretion and its consequences of black holes surviving through key epochs in cosmology [3,5] need to be studied in more details. Such investigations could constrain the initial energy fraction in black holes vis-a-vis the size of the extra dimension.

[1] For a recent review, see, D. Langlois, gr-qc/0207047 and references therein.

- [2] Ya. B. Zeldovich and I. D. Novikov, *Astron. Zh.* **43**, 758 (1966); S. W. Hawking, *Mon. Not. Roy. Ast. Soc.* **152**, 75 (1971); B. J. Carr and S. W. Hawking, *Mon. Not. Roy. Ast. Soc.* **168**, 399 (1974); S. W. Hawking, I. G. Moss and J. M. Stewart, *Phys. Rev. D* **26**, 2681 (1982); H. Kodama, M. Sasaki and K. Sato, *Prog. Theor. Phys.* **68**, 1979 (1982).
- [3] See, for instance, J. D. Barrow, E. J. Copeland, E. W. Kolb and A. R. Liddle, *Phys. Rev. D* **43**, 984 (1991); A. R. Liddle and A. M. Green, *Phys. Rept.* **307**, 125 (1998); J. Yokoyama, *Phys. Rev. D* **58**, 107502 (1998); T. Banks and W. Fischler, *hep-th/0111142*; P. He and L-Z. Fang, *Ap. J.* **568**, L1 (2002);
- [4] A. S. Majumdar, P. Das Gupta and R. P. Saxena, *Int. J. Mod. Phys. D* **4**, 517 (1995); P. S. Custodio and J. E. Horvath, *Phys. Rev. D* **58**, 023504 (1998); *Phys. Rev. D* **60**, 083002 (1999).
- [5] P.S. Custodio and J.E. Horvath, *Phys. Rev. D* **65**, 024023 (2002); D. Polarski, *Phys. Lett. B* **528**, 193 (2002); D. Blais, C. Kiefer and D. Polarski, *Phys. Lett. B* **535**, 11 (2002); R. Bean and J. Magueijo, *Phys. Rev. D* **66**, 063505 (2002); E. V. Bugaev and K. V. Konishchev, *Phys. Rev. D* **66**, 084004 (2002); L. A. Anchordoqui, J. L. Feng, H. Goldberg and A. D. Shapere, *hep-ph/0207139*.
- [6] P. Argyres, S. Dimopoulos and J. March-Russel, *Phys. Lett. B* **441**, 96 (1998); N. Dadhich, R. Maartens, P. Papadopoulos and V. Rezanian, *Phys. Lett. B* **487**, 1 (2000); R. Emparan, G. T. Horowitz, and R. C. Myers, *JHEP* **0001**, 007 (2000); R. Emparan, G. T. Horowitz, and R. C. Myers, *Phys. Rev. Lett.* **85**, 499 (2000); R. Casadio and B. Harms, *Phys. Rev. D* **64**, 024016 (2001); S. Hemming and E. Keski-Vakkuri, *Phys. Rev. D* **64**, 044006 (2001); N. Dadhich and S. G. Ghosh, *Phys. Lett. B* **518**, 1 (2001); M. Bruni, C. Germani and R. Maartens, *Phys. Rev. Lett.* **87**, 231302 (2001); P. Suranyi and L. C. R. Wijewardhana, *hep-th/0207084*.
- [7] R. Guedens, D. Clancy and A. R. Liddle, *Phys. Rev. D* **66**, 043513 (2002).
- [8] L. Randall and R. Sundrum, *Phys. Rev. Lett.* **83**, 4690 (1999).
- [9] A. R. Liddle and D. H. Lyth, *Cosmological Inflation and Large-Scale Structure*, Cambridge University Press, Cambridge (2000).
- [10] J.C. Niemeyer and K. Jedamzik, *Phys. Rev. D* **59**, 124013 (1999); A. M. Green and A. R. Liddle, *Phys. Rev. D* **60**, 063509 (1999); M. Shibata and M. Sasaki, *Phys. Rev. D* **60**, 084002 (1999); Graham D. Kribs, Adam K. Leibovich and I.Z. Rothstein, *Phys. Rev. D* **60**, 103510 (1999); T. Bringmann, C. Kiefer, D. Polarski, *Phys. Rev. D* **65**, 024008 (2002).
- [11] C. D. Hoyle, U. Schmidt, B. R. Heckel, E. G. Adelberger, J. H. Gundlach, D. J. Kapner and H. E. Swanson, *Phys. Rev. Lett.* **86**, 1418 (2001).